



Effect of Falgwadi Yoga on Changes in Biochemical Markers of Iron Deficiency Anaemia

Shah Rahul¹, Pani S.¹, Arora M. T.², Gujarathi R. H.³

¹Department of Kaumarabhritya, Parul Institute of Ayurveda, Parul University, Vadodara, Gujarat, India

²Department of Swasthavritta, Government Ayurved College, Vadodara, Gujarat, India

³Department of Kaumarabhritya, Bharati Vidyapeeth (DU), College of Ayurved, Pune, Maharashtra, India

Correspondence Email: shrigujarathi@gmail.com

ABSTRACT

Iron deficiency is by far the first cause of nutritional anaemia worldwide. A large portion of iron deficiency is preventable with appropriate and timely intervention. Anaemia has been a prevalent health issue in India with over 70% children. Nearly 73 million children below the age of three (79%) suffer from varying degrees of anaemia. Pāṇḍu is supposed to be the nearest co-related entity to anaemia. The management of this syndrome is supplementing elemental iron or bio-iron through food or as medicine. Elemental iron supplements have their own drawbacks and present as unwarranted effects like constipation, vomiting, metallic taste, non-acceptance and compliance, iron deposition in the tissues and sometimes iron toxicity. Bioiron i.e. iron of natural origin (plant and animal origin) bypasses these symptoms, it is readily absorbed and readily bio-converted. The present study supplements bio iron i.e. iron of natural origin using Santarpana Cikitsa with Falgu (Añjira), Kṛṣṇa Drākṣa (Manukā) and Kharjura combination was undertaken in the management of Pāṇḍu (iron deficiency anaemia) in children. The effect of these supplements were studied on anaemia markers and it was observed that it had positive effect on rise in haemoglobin, serum iron level and serum ferritin levels with a marked fall in total iron binding capacity, on administration of the trial drug for 30 days.

Keywords: Iron Deficiency Anemia, Hemoglobin, Serum Ferritin, Serum Iron, Figs, Dates, Black Currants

Received 09.11.2022

Revised 20.01.2023

Accepted 25.01.2023

INTRODUCTION

Anaemia has been a prevalent health issue in India. According to the National Family Health Survey (NFHS) data for anaemia prevalence among children under three years of age shows 79% which is five percent more than the previous survey conducted. Nearly 73 million children below the age of three (79%) suffer from varying degrees of anaemia. Over 50million children suffer from moderate-to-severe anaemia. According to NFHS study, the highest prevalence is among the six to twenty-three months of age group. Based on studies by the National Nutrition Monitoring Bureau, anaemia prevalence among children between one to five years of age is around 66%, with varying figures of 33 to 93% across different states of India. There is an irreparable damage that anaemia in childhood can cause particularly to the development of a young child, physically and intellectually. The numbers are staggering, about two billion people—over30% of the world's population—are anaemic, mainly due to iron deficiency.

IDA can be understood as a blood disorder in which the blood has a reduced ability to carry oxygen due to a lower than normal number of red blood cells, or a reduction in the amount of hemoglobin. Anaemia is the most common disorder of the blood, of the developing world. "Anaemia" is not a diagnosis in itself. It is an objective sign of the presence of other diseases. iron deficiency is considered to be the most common cause of anaemia, at least in children and females, though other conditions of nutrition such as vitamin B₁₂ and folate deficiencies, parasitic infections, chronic gut inflammation leading to reduced absorption of iron, chronic diseases, endocrine disorder and various inherited disorder can also causes anaemia [1, 2].

Table 1: W.H.O. haemoglobin thresholds used to define anaemia [1 gm/dl:0.6206mmol/L] [3]

Age/Gender Group	Hb Threshold [gm/dl]	Hb Threshold [mmol/L]
Children (0.5–5.0 yrs)	11.0	6.8
Children (5–12 yrs)	11.5	7.1
Teens (12–15 yrs)	12.0	7.4
Women, non-pregnant (>15yrs)	12.0	7.4
Men (>15yrs)	13.0	8.1

0.5 g of iron is present in iron a full-term new-born as against of 5 g in adults. An average of 0.8 mg of iron needs to be absorbed daily during the initial 15 years of life [4, 5]. Small additional amount of iron is necessary to overcome the normal losses of iron caused by shedding of cells. Thus it necessitates absorption of about 1mg of iron daily to maintain positive iron balance during childhood. Less than 10% of dietary iron is usually absorbed, and thus 8-10 mg of iron is necessary daily, to maintain iron levels.

During infancy, when growth is rapid, iron from cow's and breast milk makes it difficult to maintain body iron levels at the optimum. Breastfed infants have are at an advantage, because they can absorb iron 2-3 times more efficiently than infants that are fed cow's milk. Breastfed infants are at risk to land in to iron deficiency if regular intake of iron-fortified foods by six months of age is not started [6]. The usual dietary pattern observed in infants and toddlers with nutritional iron-deficiency anemia, is excessive consumption of cow's milk [7, 8, 9] (low iron content, blood loss from milk protein colitis). Under nutrition is also generally responsible for iron deficiency.

Iron deficiency and iron-deficiency anaemia

Iron deficiency is a condition where serum iron levels fall below the threshold levels i.e. 50 mg/dL but the pathophysiology of anaemia has not set off. Iron deficiency has non hematologic systemic effects. Both iron deficiency and iron-deficiency anaemia are associated with impaired neurocognitive function in infancy [9]. Iron-deficiency anaemia may also be associated with, possibly irreversible, cognitive defects [10]. Some studies suggest an increased risk of seizures, strokes, breath-holding spells in children, and exacerbations of restless legs syndrome [11]. Due to the potential of adverse neurodevelopmental outcomes, reducing the frequency of iron deficiency has to be an important goal, overall. Other non-hematologic consequences of iron deficiency include pica and pagophagia, the desire to ingest ice.

Diagnosis of IDA [12]

A thorough medical history is important to the diagnosis of iron-deficiency anemia. A travel history to areas in which hookworms and whipworms are endemic may also be helpful in guiding certain stool tests for parasites or their eggs [13]. In progressive iron deficiency, sequential biochemical and hematologic events occur. Tissue iron stores deplete. The reduced serum ferritin reflects this depletion, Ferritin is an iron-storage protein, which can provide an estimate of body iron stores when inflammatory diseases are absent [14]. Later serum iron levels decrease. The iron-binding capacity of the serum (serum transferrin) increases, and the transferrin saturation falls below normal. As iron stores decrease, iron becomes unavailable to form a complex with protoporphyrin to form haem. At this point, iron deficiency progresses to iron-deficiency anemia [15]. With lesser hemoglobin, the red blood cells (RBCs) become smaller called microcytes, and varied in size. The changes are associated with a decrease in mean corpuscular volume (MCV) and mean corpuscular hemoglobin [16]. The RBC count decreases. The reticulocyte count (percentage) is normal or moderately elevated. The peripheral blood smear reveals hypochromic, microcytic RBCs with substantial variation in cell size. A presumption of iron-deficiency anemia is most of the times done by a complete blood count (CBC) showing a microcytic anemia picture with high RDW, decreased RBC count (Hematocrit), normal WBC count, and a normal or elevated platelet count. Other laboratory investigations like the reduced serum ferritin, reduced serum iron, and increased total iron-binding capacity, generally, are not necessary unless severe anemia requires a faster diagnosis, other complicating clinical signs are present, or the anemia does not respond to oral iron therapy. A diagnosis of iron deficiency in the absence of anemia is challenging. Serum ferritin is a useful tool, whose value is increased by also measuring C-reactive protein to help identify false-negative results because of concomitant inflammation.

Table 2: Laboratory studies differentiating diagnosis of microcytic anaemia [16, 17]

Study	Iron Deficiency Anaemia	A or β Thalassemia	Anaemia of Chronic Disease
Haemoglobin	Decreased	Decreased	Decreased
Mean Corpuscular Volume	Decreased	Decreased	Normal / Decreased
RDW	Increased	Normal / Minimally Increased	Normal / Increased
RBC	Decreased	Normal / Increased	Normal / Decreased
Serum Ferritin	Decreased	Normal	Increased
TIBC	Increased	Normal	Decreased

Serum Transferrin Saturation	Decreased	Normal	Decreased
FEP	Increased	Normal	Increased
Soluble Transferrin Receptor	Increased	Normal	Normal
Reticulocyte Haemoglobin Concentration	Decreased	Normal	Normal / Decreased

The response of iron-deficiency anaemia to adequate administration of iron is a critical diagnostic and therapeutic feature. Oral administration ferrous salts [18, 19] most often ferrous sulphate is inexpensive and effective. No evidence is available regarding the addition of any trace metal, vitamin, or other hematinic substance leading to significant increase in the response to simple ferrous salts. Calcium and dietary fibre decrease the absorption of iron, but this can be overcome by co-administration of ascorbic acid. Tea is a significant inhibitor of iron absorption. The therapeutic dose should be calculated in terms of elemental iron. Iron in a total daily dose of 3-6 mg/kg of elemental iron in 1 or 2 doses is enough, while higher doses are used in severe cases. Parenteral iron is considered in malabsorption, where compliance is inadequate, as oral therapy is otherwise effective, less expensive and safe. Iron therapy increases the virulence of malaria and certain gram-negative bacteria. Iron overdose is mostly associated with Yersinia infection. Dietary counselling is usually necessary. Intake of cow's milk, needs be limited. Dietary iron should be increased. Iron from haem sources is more bioavailable than from non-haem sources. Iron should be continued for 2-3 months after values normalize so as to re-establish iron stores.

The dried fruits of *Fig* are reported to be an important source of different vitamins, minerals, carbohydrates, sugars, organic compounds, phenolic etc. [20-21]. The fresh as well as dried fruits of figs are good source of fibre and polyphenols [22-23]. Figs are quoted in the traditional systems of medicine like as *Āyurveda*, *Unānī*, and *Siddha* and are used to treat diseases of the endocrine system (diabetes), respiratory system, digestive tract (indigestion and vomiting), menstrual disorders and menstrual pain etc. As per data of USDA for the mission variety, dried figs are rich in fibre, copper, manganese, magnesium, potassium, calcium, iron and vitamin K. Fruits also have other nutrients in smaller proportion. Fruits of *Ficus carica* (*Figs*) are consumed fresh or dried, which are an excellent source of minerals, vitamins, carbohydrates, and dietary fibre, having a low fat and cholesterol content and ample amount of amino acids [24-26].

Kharjūra term is used for fruits of Phoenix dactylifera Linn. Dates fruits are rich in carbohydrates in the form of sucrose, glucose and fructose (65% – 80%), and are a good source of fibres and certain essential minerals. *Kharjūra* or dates fruits contain low fats and protein but contain high sugar concentration and do not have starch. *Kharjūra* or dates fruits are rich source of niacin, B6, riboflavin, thiamine, pantothenic acid, manganese, magnesium, iron and phosphorus. *Kharjūra* or dates fruits also contain a good amount of zinc, copper, potassium selenium and folic acid.

Dried fruits of Grape Vine (*Vitis Vinifera*) i.e Black Currant is considered as *Drākṣā* in *Āyurveda*. The important basic constituent of grape is manganese. *Vitis Vinifera* or grape vine contains vitamin B6, thiamine, riboflavin, vitamin C and potassium. Various studies concluded that it also contains a chemical substance like resveratrol, a polyphenol having antioxidant properties [27].

Table 3: Nutritional analysis

	<i>Añjira</i>	<i>Drākṣā</i>	<i>Kharjūra</i>
Edible Portion (Percentage)	99.0	98.0	94.0
Moisture (Grams)	38.0	20.2	15.3
Protein (Grams)	1.3	0.3	2.5
Total Fat (Grams)	0.2	0.5	0.4
Total Minerals (Grams)	0.6	1.2	3.15
Total Edible Fibre (Grams)	2.2	1.7	3.9
Carbohydrates (Grams)	57.6	75.2	75.8
Total Calories (Kcal)	373	316	317
Calcium (Milligrams)	80.0	130.0	120.0
Phosphorus (Milligrams)	30.0	110.0	50.0
Iron Compounds (Grams)	0.87	0.69	1.54
Elemental Iron (Milligrams)	102	7.7	173
Carotene (Micrograms)	162.0	21.0	26.0
Thiamine (Milligrams)	0.06	0.03	0.01
Riboflavin (Milligrams)	0.05	0.14	0.02
Niacin (Milligrams)	0.6	0.4	0.9
Ascorbic Acid (Milligrams)	5.0	1.0	3.0
Essential Amino Acids (Total Nitrogen Gram Percent)	2.8	2.3	0.4

MATERIAL AND METHODS

An interventional study was conducted in an urban and semi urban area at two centers simultaneously. Ethical clearances were obtained from both the centers. The trial was registered with CTRI (Central Trial Registry of India) and a trial number was obtained on 22.10.2019, bearing no. CTRI/2019/10/021749. The patients were asked to consume a freshly prepared paste containing *Falgu (Anjira)*, *KṛṣṇaDrākṣā (Manukā)* and *Kharjūra*, 10 grams each, soaked overnight in water, for the span of 30 days. A biochemical examination for assessment of a complete blood count, serum iron, serum ferritin and total iron binding capacity was done before the commencement and after the end of the intervention. The pre commencement investigations were done for confirmation of diagnosis of IDA and post-trial were done to evaluate the effect of intervention.

RESULTS AND DISCUSSION

The data collected in the present study in respect of biochemical parameters are detailed below and were studied for improvement in individual factors and the results were subjected to statistical analysis.

Table 4: Effect of interventional combination on anaemia profile parameters

Parameter	Haemoglobin	Serum Iron	Serum Ferritin	TIBC
Mean BT	7.9475	35.063	17.375	535.88
Mean AT	10.056	70.8	39.713	339.25
X	2.1088	35.738	22.338	196.63
% of improvement	20.97%	50.48%	56.25%	57.96%
't' value	-29.57	-42.25	-19.37	20.337

The mean grade of Haemoglobin on 1st day was 7.94 which increased to 10.056 at 30th day. The mean increment in score was 20.97% which is significant as observed by paired t test (as p value <0.05) thus it can be said that there is significant increment on Haemoglobin. Increase in haemoglobin is not only dependent on supply of iron; though iron has a major role to play in rise in haemoglobin, but also on supply on other nutrients as the molecule is made up of two contents; the haem and the globin. Dietary iron also has an influence on the rise in haemoglobin [21-24].

The mean grade of Serum Iron on 1st day was 35.06 which increased to 70.8 at 30th day. The mean increment in score was 50.48% which is significant as observed by paired t test (as p value <0.05) thus it can be said that there is significant increment on Serum Iron. This probably might be due to better absorption of iron from the interventional combination as it was a biological form of iron [25,26].

The mean grade of Serum Ferritin on 1st day was 17.37 which were increased to 39.71 at 30th day. The mean increment in score was 56.25% which is significant as observed by paired t test (as p value <0.05) thus it can be said that there is significant increment on Ferritin. Ferritin also rises during inflammatory conditions and so is considered as an inflammatory marker. In this study, inflammation was not ruled out on serological parameters, and so, it is difficult to comment whether there were any inflammatory conditions present in the subjects post completion of study, though visually there were none nor there were any conditions reported by the study subjects to suspect so [27].

The mean grade of TIBC on 1st day was 535.88 which decreased to 339.25 at 30th day. The mean increment in score was 57.96% which is significant as observed by paired t test (as p value <0.05) thus it can be said that there is significant fall in TIBC. The fall in Total Iron Binding Capacity was consistent with the findings of rise in haemoglobin levels and serum iron levels. Total Iron Binding Capacity is inversely proportional to the serum iron levels and haemoglobin levels.

CONCLUSION

The increment in haemoglobin and serum iron level concentration and total iron binding capacity is significant, though the haemoglobin has not reached the normal levels in the 30 days span. The increment of levels of ferritin is better. The trial combination does not give rise to any adverse/untoward/unwarranted reactions of the drug as are commonly seen with the commonly used iron salts.

REFERENCES

1. Smith Jr, R. E. (2010). The clinical and economic burden of anemia. *Am J Manag Care*, 16(Suppl Issues), S59-S66.
2. Vos, T., Barber, R. M., Bell, B., Bertozzi-Villa, A., Biryukov, S., Bolliger, I., & Brugha, T. S. (2015). Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The lancet*, 386(9995), 743-800.
3. Zimmermann, M. B., & Hurrell, R. F. (2007). Nutritional iron deficiency. *The lancet*, 370(9586), 511-520.

4. Ashish, K. C., Rana, N., Målqvist, M., Ranneberg, L. J., Subedi, K., & Andersson, O. (2017). Effects of delayed umbilical cord clamping vs early clamping on anemia in infants at 8 and 12 months: a randomized clinical trial. *JAMA pediatrics*, *171*(3), 264-270.
5. Baker, R. D., Greer, F. R., & Committee on Nutrition. (2010). Diagnosis and prevention of iron deficiency and iron-deficiency anemia in infants and young children (0–3 years of age). *Pediatrics*, *126*(5), 1040-1050.
6. Berglund, S. K., Westrup, B., & Domellöf, M. (2015). Iron supplementation until 6 months protects marginally low-birth-weight infants from iron deficiency during their first year of life. *Journal of pediatric gastroenterology and nutrition*, *60*(3), 390-395.
7. Lai, F. P., & Yang, Y. J. (2018). The prevalence and characteristics of cow's milk protein allergy in infants and young children with iron deficiency anemia. *Pediatrics & Neonatology*, *59*(1), 48-52.
8. Camashella, C. (1832). Iron Anemia Nejm. *The New England Journal of Medicine*.
9. Congdon, E. L., Westerlund, A., Algarin, C. R., Peirano, P. D., Gregas, M., Lozoff, B., & Nelson, C. A. (2012). Iron deficiency in infancy is associated with altered neural correlates of recognition memory at 10 years. *The Journal of pediatrics*, *160*(6), 1027-1033.
10. Falkingham M, Abdelhamid A, Curtis P, et al. The effects of oral iron supplementation on cognition in older children and adults: a systematic review and meta-analysis. *Nutr J*. 2010;9:4
11. Powers, J. M., & Buchanan, G. R. (2017). Potential for improved screening, diagnosis, and treatment for iron deficiency and iron deficiency anemia in young children. *The Journal of pediatrics*, *188*, 8-10.
12. Peyrin-Biroulet, L., Williet, N., & Cacoub, P. (2015). Guidelines on the diagnosis and treatment of iron deficiency across indications: a systematic review. *The American journal of clinical nutrition*, *102*(6), 1585-1594.
13. Jourdan, P. M., Lambertson, P. H., Fenwick, A., & Addiss, D. G. (2018). Soil-transmitted helminth infections. *The Lancet*, *391*(10117), 252-265.
14. Abdullah, K., Birken, C. S., Maguire, J. L., Fehlings, D., Hanley, A. J., Thorpe, K. E., & Parkin, P. C. (2017). Re-evaluation of serum ferritin cut-off values for the diagnosis of iron deficiency in children aged 12-36 months. *The Journal of Pediatrics*, *188*, 287-290.
15. Centers for Disease Control and Prevention (CDC). (2002). Iron deficiency--United States, 1999-2000. *MMWR. Morbidity and mortality weekly report*, *51*(40), 897-899.
16. Champion, E. W., & Deloughery, T. G. (2014). Microcytic anemia. *N Engl J Med*, *371*, 1324-1331.
17. Ferrara, M., Capozzi, L., Russo, R., Bertocco, F., & Ferrara, D. (2010). Reliability of red blood cell indices and formulas to discriminate between β thalassemia trait and iron deficiency in children. *Hematology*, *15*(2), 112-115.
18. McDonagh, M. S., Blazina, I., Dana, T., Cantor, A., & Bougatsos, C. (2015). Screening and routine supplementation for iron deficiency anemia: a systematic review. *Pediatrics*, *135*(4), 723-733.
19. Powers, J. M., & Buchanan, G. R. (2014). Diagnosis and management of iron deficiency anemia. *Hematology/Oncology Clinics*, *28*(4), 729-745.
20. Jeong, W. S., & Lachance, P. A. (2001). Phytosterols and fatty acids in fig (*Ficus carica*, var. Mission) fruit and tree components. *Journal of food science*, *66*(2), 278-281.
21. Veberic, R., Colaric, M., & Stampar, F. (2008). Phenolic acids and flavonoids of fig fruit (*Ficus carica* L.) in the northern Mediterranean region. *Food chemistry*, *106*(1), 153-157.
22. Vinson, J. A., Zubik, L., Bose, P., Samman, N., & Proch, J. (2005). Dried fruits: excellent in vitro and in vivo antioxidants. *Journal of the American College of nutrition*, *24*(1), 44-50.
23. Vinson, J. A. (1999). The functional food properties of figs. *Cereal foods world*, *44*(2), 82-87.
24. Slatnar, A., Klancar, U., Stampar, F., & Veberic, R. (2011). Effect of drying of figs (*Ficus carica* L.) on the contents of sugars, organic acids, and phenolic compounds. *Journal of Agricultural and Food Chemistry*, *59*(21), 11696-11702.
25. Veberic, R., Colaric, M., & Stampar, F. (2008). Phenolic acids and flavonoids of fig fruit (*Ficus carica* L.) in the northern Mediterranean region. *Food chemistry*, *106*(1), 153-157.
26. Solomon, A., Golubowicz, S., Yablłowicz, Z., Grossman, S., Bergman, M., Gottlieb, H. E., & Flaishman, M. A. (2006). Antioxidant activities and anthocyanin content of fresh fruits of common fig (*Ficus carica* L.). *Journal of agricultural and food chemistry*, *54*(20), 7717-7723.
27. Iriti, M., & Faoro, F. (2006). Grape phytochemicals: a bouquet of old and new nutraceuticals for human health. *Medical hypotheses*, *67*(4), 833-838.

CITATION OF THIS ARTICLE

Shah, Rahul, Pani, S., Arora, M. T., Gujarathi, R. H. Effect of falgwadi yoga on changes in biochemical markers of iron deficiency anaemia. *Bull. Env. Pharmacol. Life Sci.*, Vol 12[3] Feb 2023 : 170-174.