



**ORIGINAL ARTICLE**

**OPEN ACCESS**

## **Fractionation and Purification of Bioactive Peptides in Excretory/Secretory Products of Third Instar Larvae of *Chrysomya Megacephala* (Calliphoridae: Diptera)**

**Nancy Taha Mohamed**

Lecturer of Entomology, Zoology & Entomology Department, Faculty of Science, Helwan University.

EMAIL ADDRESS: [nancyt0000@yahoo.com](mailto:nancyt0000@yahoo.com)

### **ABSTRACT**

*The beneficial effects of maggots on wounds have been attributed to various mechanisms notably the debridement (degradation) of necrotic tissue. It was demonstrated that the maggots ES exhibit antibacterial actions against both Gram-positive and Gram-negative bacteria, including MRSA, Escherichia coli and Pseudomonas aeruginosa. The present study revealed the presence of 40 protein fractions in excretory/secretory products of third instar larvae of Chrysomya megacephala, only 3 protein fractions showed antibacterial activity against Bacillus subtilis, number 22, 23 and 24. The separated protein fraction number 22 showed protein bands at 28 and 31 KDa protein, separated protein fraction number 23 showed protein bands at 17, 24 and 28 KDa and separated protein fraction number 24 showed protein band at 26 KDa.*

**Key Words:** *Chrysomya megacephala, antibacterial peptides, Excretory/Secretory products*

Received 11.08.2015

Revised 21.08.2015

Accepted 05.09.2015

### **INTRODUCTION**

Maggot therapy has been traditionally practiced for debridement of necrotic wounds as well as for curing bacterial infections at the wounds site [1]. It has been reported to have advantages over the conventional methods, especially for treatment of wounds infected by multi-drug resistant methicillin-resistant *Staphylococcus aureus* (*S. aureus*) or MRSA [2]. Their antiseptic action has been investigated by many researchers in the past for specialized antibacterial properties or presence of antimicrobial factor(s). The beneficial effects of maggots on wounds have been attributed to various mechanisms notably the debridement (degradation) of necrotic tissue. It was originally believed that this debriding action of maggots was restricted only to their mechanical wriggling, but recently many proteolytic enzyme classes have been isolated from maggot excretions and secretions (ES) which are able to specifically dissolve the laminin and fibronectin of the extracellular matrix in the necrotic tissue. This liquifies the dead tissues enabling the maggot to take it up by suction [3]. Finally, maggots promote wound healing, stimulate granulation and promote the formation of human fibroblasts [4].

In particular, the mechanism of action of maggot disinfection on wounds was studied [5]. They found that the excretion of maggots exhibited a strong and rapid disinfection action on *S. aureus*. Subsequently, several other groups collected the excretions/secretions (ES) of maggots and screened it on various microbes [6]. Many species of blowfly maggots were investigated against both gram negative and gram-positive bacteria [7]. Several recent studies have demonstrated that the maggots ES from aseptically-raised *Lucilia sericata* larvae exhibit antibacterial actions against both Gram-positive and Gram-negative bacteria, including MRSA, *Escherichia coli* and *Pseudomonas aeruginosa* [6]. In addition, maggots can ingest bacteria as part of their normal feeding process [3]. The antibacterial activity of *Lucilia cuprina* maggot ES was previously shown to have partial growth inhibition on gram-positive, *S. aureus* [8].

Recently, a low molecular weight insect peptide was found mainly responsible for the antimicrobial activity of the maggots, when exposed to an infectious environment of a wound. The peptide, lucifencin, that belongs to the insect antimicrobial peptides (AMP), defensin family, was reported to be secreted in the hemolymph, body fat and ES of the maggots. The maggots were raised in an environment

simulating a wound that increased the production of the AMP as an innate immune response of the insect [9].

*Chrysomya megacephala* (F.), the Oriental latrine fly, is a common blow fly species of medical importance in many parts of the world, including Egypt. Adults may feed on food sources including nectar, animal carcasses, garbage, and other filth materials, or even human food. Therefore, it is possible that mechanical transfer of potential disease causing pathogens, such as bacteria, viruses, protozoa, and helminthes eggs, to human food may occur [10]. Larvae of this species are known to cause myiasis in several mammal species, including humans [11]. Another facet of medical importance of this blow fly is its association with human corpses and its relevance to forensic entomology. Many researchers have reported that specimens of *C. megacephala* were found connected with cases of human death [12]. Recent molecular and biochemical studies on third instar larvae of *C. megacephala* revealed the presence of serine proteases especially trypsin and chymotrypsin in excretory/secretory products [13 , 14 ].Also the excretory/secretory products of third instar larvae of *Chrysomya megacephala* showed antibacterial activity against *Escherichia coli*, *Staphylococcus aureus* and *Bacillus subtilis* with nearly more activity against *Bacillus subtilis* followed by *Staphylococcus aureus* and *Escherichia coli* [14 ] .

The aim of the present study is to fractionate the excretory/secretory products of third instar larvae of *Chrysomya megacephala*, to screen these protein fractions against *Bacillus subtilis* (as ES products has more activity against it) and finally to purify different peptides from the active protein fractions in order to clarify the usefulness of ES from *Chrysomya megacephala* in maggot therapy.

## **MATERIALS AND METHODS:**

### **REARING OF INSECT:**

The laboratory colony of *C. megacephala* used in this study was established in the Department of Entomology, Faculty of Science, and Helwan University. *C. megacephala* was reared according to following protocol [15]. They were identified according to reported method [16]. Adults from the stock colony of *C. megacephala* were kept in cages (38×38×56 cm) at 25±3°C, 14h photoperiod and 60–70% R.H. The cages were made with a wooden floor, a glass roof, and wire gauze on three of the sides. The fourth side was wooden with a circular hole fitted with a cloth sleeve to facilitate daily feeding, cleaning of the cage, and removal of eggs. Adults were supplied daily with granular sucrose, water, and pieces of liver.

Water was supplied by dipping a piece of cotton as a wick in a bottle filled with water, and the liver was provided in a Petri- dish. Egg batches were removed daily and transferred to a fresh piece of chicken placed in a rearing enamel bowl (35 cm in diameter) covered with muslin secured with a rubber band. At the prepupal stage, dry autoclaved sawdust was added to the bowl as a medium for pupation. Pupae were sieved from the sawdust and transferred to adult cages described above for adult emergence.

### **COLLECTION OF LARVAL SECRETIONS:**

The ES products were collected according to modifications of reported method [17]. Native excretions/secretions (nES) were collected by incubating third instars larvae of *C. megacephala* in (50,000 larvae/ 500 ml) sterile distilled water for 1h at 30°C in darkness. The sterile liquid was siphoned from the containers and centrifuged at 10,000×g for 5 min to remove particulate material, after which the supernatant was collected and separated.

### **FRACTIONATION OF NATIVE EXCRETIONS/SECRETIONS (NES):**

Gel filtration technique was used to purify antimicrobial active compounds from the crude sample investigated. The Pharmacia column (40×2 cm) was packed by Sephadex G 150 after its soaking with phosphate buffer (pH 7.2) overnight and the samples were mixed with blue dextran and bromophenol blue dyes, loaded on the top of the column and fractions were collected (5 ml of each fraction), fractionation was performed using LPLC system (pump model MINI Puls3, detector UVVIS-151 FC 203B, Gilson).

### **SCREENING ANTIBACTERIAL ACTIVITY OF PROTEIN FRACTIONS**

#### **AGAR WELL DIFFUSION METHOD**

The antibacterial activity of protein fractions was determined using agar well diffusion method [19]. All the protein fractions were tested *in vitro* for their antibacterial activity against Gram positive bacteria *Bacillus subtilis* using nutrient agar medium. Ampicillin was used as standard drugs for *Bacillus subtilis*. DMSO was used as solvent control. The protein fractions were tested at a concentration of 1 mg/ml against the bacterial strain.

The sterilized media was poured onto the sterilized Petri dishes (20-25 ml, each petri dish) and allowed to solidify. Wells of 6 mm diameter was made in the solidified media with the help of sterile borer. A sterile swab was used to evenly distribute bacterial suspension over the surface of solidified media and solutions of the test compounds were added to each well with the help of micropipette. The plates were

incubated at 37°C for 24 hrs for antibacterial activity. This experiment was carried out in triplicate and zones of inhibition were measured in mm. scale.

#### **SODIUM DODECYL SULPHATE (SDS- PAGE) FOR ANTIMICROBIAL PEPTIDES**

Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis (SDS-PAGE) was used to Separation of Proteins according to their molecular weight.

#### **Reagent and buffers:**

Acrylamide Monomer Stock Solution (30%): 29.2 gm acrylamide and 0.8 gm N,N'-methylene bis-acrylamide were dissolved in 100 ml distilled water. (**Resolving Buffer:** 1.5M Tris-HCl, pH 8.8.;**Stacking Buffer:** 0.5M Tris-HCl, pH 6.8., **Resolution Buffer,** pH 8.3: 0.192 M Glycine, 0.02 Trisbase and 0.5 % SDS, **Sample Buffer:** 12 ml 1.0 m Tris-HCL pH 6.8, 8 ml glycerol, 4 ml 2-mercaptoethanol, 16 ml 10% SDS and 2 ml 0.05% bromophenol blue were made up to 80 ml with distilled water).

#### **Preparation of resolving gel (10%)**

The gel was prepared from: 4.0 ml distilled water, 2.5 ml resolving buffer, 3.7 ml acrylamide monomer, 100 µl 10% SDS, 50 µl 10% ammonium persulphate and 5 µl Tetramethylethylene Diamine (TEMED).

The gel was poured between the glass plates immediately and carefully overlaid with a layer of distilled water, and then the gel was kept at room temperature to polymerize.

#### **Preparation of stacking gel (4%)**

The gel was prepared from: 3.06 ml distilled water, 2.5 ml stacking buffer, 0.65 ml acrylamide monomer, 100 µl 10% SDS, 50 µl 10% ammonium persulphate and 5 µl TEMED. The surface of the polymerized resolving gel was rinsed with 5-7 mL of stacking gel solution. The comb was aligned in the proper position, then, the stacking gel solution was added up to 2 mm from the top edge of the resolving gel, then left for polymerization at room temperature.

#### **Sample preparation:**

30 µl of each active fraction were mixed with 10 µl of loading sample dye. The samples were then boiled in a water bath for 2 minutes. The samples were then applied to the gel wells.

#### **Running Conditions:**

Electrophoresis was carried out with constant volt of 60V, The run was terminated when the bromophenol blue marker reached to the bottom of the gel. The separated proteins on polyacrylamide gels were then stained with Coomassie blue R-250 according to the method described by Merrill ( 18).

The electrophoreses gel was soaked in excess of staining solution for two hours, then the gel was rinsed with distilled water and distained with excess amount of distaining solution for several times until the excess stain was removed.

## **RESULTS**

The present study revealed the presence of 40 protein fractions in excretory/secretory products of third instar larvae of *Chrysomya megacephala*, only 3 protein fractions showed antibacterial activity against *Bacillus subtilis*, number 22,23 and 24(Table 1). The separated protein fraction number 22 showed protein bands at 28 and 31 KDa protein(Fig.1 ,Lane 1), separated protein fraction number 23 showed protein bands at 17 , 24 and 28 KDa (Fig.1 ,Lane 2) and separated protein fraction number 24 showed protein band at 26 KDa (Fig.1 ,Lane 3).


Table (1):Mean zone of inhibition in mm ± Standard deviation against *Bacillus subtilis* using different fractions. Results are depicted in the following table:

<b>Fraction number</b>	<b>Mean zone of inhibition in mm ± Standard deviation</b>
<b>22</b>	<b>21.3 ±1.2</b>
<b>23</b>	<b>20.1± 1.5</b>
<b>24</b>	<b>17.6± 0.58</b>

## **DISCUSSION**

The present study revealed the presence of 40 protein fractions in excretory/secretory products of third instar larvae of *Chrysomya megacephala*, only 3 protein fractions showed antibacterial activity. On purification of these protein fractions, they showed protein bands between 17KDa and 31 KDa.

Since 2000, several research groups have been aiming to isolate and characterize such antimicrobial peptides from the ES by utilizing current methods of protein purification. In the laboratory of [19] the ES of maggots was fractionated using an ultra filtration device with a 10 KDa and 500 Da molecular weight cut-off membrane generating three fractions of molecular weights: >10 kDa,500 Da-10 KDa and <500 Da.



**Fig : (1):** SDS-PAGE of separated proteins compared with standard molecular weight proteins, protein fraction number 22(Lane 1),protein fraction number 23 (Lane 2),protein fraction number 24 (Lane 3),Marker molecular weight (Lane 4).

The activity against *S. aureus* was detected in <500 Da fraction and 500 Da–10 KDa fraction, but not in the fraction above 10 KDa. Even though these fractions were investigated in further detail regarding their physicochemical properties and antimicrobial activities [19], their constituents were not identified. The antimicrobial properties of *L. sericata* larval ES and the attempts to characterize its components were independently studied in several other laboratories [6]. For example, the study of Kerridge *et al.* [20] revealed in the secretions the presence of small (<1 kDa) antimicrobial factors active against Gram-positive bacteria such as *S. aureus*, including both methicillin-resistant *S. aureus* (MRSA) and methicillin-sensitive *S. aureus* (MSSA), and *Streptococcus pyogenes*. However, Gram-negative *Pseudomonas aeruginosa* was less sensitive. This active factor passed through the filter of the 3 kDa cut-off when the secretion was fractionated by ultrafiltration procedure. In this case, anti-MRSA activity was also detected in the retentates of the 10 kDa and 5 kDa filters indicating the presence of at least one additional larger antimicrobial agent. The authors concluded that the activities in the secretions possess characteristics consistent with insect antimicrobial peptides and are considered to be of low molecular weight, highly stable and a systemic part of the larva [20].

Jana *et al.*, [21] partially purified a compound of maggot ES of *Lucilia sericata* with antibiofilm properties against *S. aureus* isolate with a size of around 25 KDa. This molecular weight is similar to those separated in the present study and this may indicate that these purified peptides from maggots ES of third instar larvae of *C. megacephala* with antibiofilm properties against *S. aureus*.

Insect immunity system is based on the recognition of microbial cell wall components, such as lipopolysaccharide (Gram-negative bacteria), peptidoglycans and lipoteichoic acids (Gram-positive bacteria), as well as  $\alpha$ -1,3-glucans (fungi) by specific receptors. Insects' defense mechanisms generally are divided into two categories: non-specific and specific (innate immune system). In non-specific defense, wide varieties of mechanisms: including morphological, behavioral, developmental (like immunity at maturity), physiological, nutritional, biochemical and genetic are being rendered [22]. Specific mechanisms, which described above, are subdivided into cellular and humoral immune systems [23]. The humoral immune system, is based on the production of induced antimicrobial peptides (AMPs) against bacterial and fungal pathogens [23]. AMPs are synthesized in insect fat body and or in some blood cells, and secreted into haemolymph [24]. Antimicrobial peptides have mainly antifungal and antibacterial properties [25]. In insects, the constitutive AMPs are accumulated in blood cells and salivary glands. Upon a microbial challenge the AMPs get released into the haemolymph [26]. In contrast, induced AMPs are synthesized in fat bodies right after the microbial infection takes place and subsequent to their biosynthesis, AMPs release into the blood stream [27]. Meanwhile, it is noteworthy that other organs and tissues such as haemocytes might be involved in AMPs synthesis [28]. Insect antimicrobial peptides are classified into four major groups; cationic peptides, anionic peptides, aromatic dipeptides and also peptides derived from oxygen-binding proteins [29]. Most found AMPs are classified as cationic peptides with 12-50 amino acids in length [30].

Cationic peptides bear 30-50% hydrophobic amino acid residues. These peptides have a wide range of activities such as antibacterial, antifungal, antiviral (AntiHIV, AntiHSV), anticancer, antiparasitic as well as antiendotoxin activities [30].

Cationic peptides can be subdivided into four broad classes on the basis of biochemical and structural features; a) linear  $\alpha$ -helical peptides, b) cyclic peptides containing one or several disulfide bridges, c) proline and glycine-rich peptides, d)  $\beta$ -sheets peptides [30].

Of mode of action, proline-rich peptides are more active against Gram-negative bacteria than Gram-positive bacteria. The higher resistance of Gram-positive bacteria might be due to the peptide decomposition by activity of bacterial extracellular proteases. Glycine-rich peptides have molecular weight between 8-30kDa. These are mostly active against Gram-negative bacteria and including Attacin, Gloverin, Dipterin, and others [31]. Although the peptides purified from ES products of third instar larvae of *C. megacephala* may be cationic peptides as the range of peptides purified in the present study is from 17kDa to 31kDa, but they are active against gram positive as *B.stibulis* and *S.aureus*. This may be due to that peptides purified from maggot ES of third instar larvae of *C. megacephala* are not decomposed by activity of bacterial extracellular proteases.

In general, mechanisms of action of antimicrobial peptides are mainly depended on both the molecular characteristics of the peptide and the metabolic states of the microbial cells [32]. Although mechanism of action of AMPs is a controversial subject, but the general consensus is that the majority of antimicrobial peptides, if not all, remarkably disrupt the cell membranes by making pores or channels, leading to the release of ions that may disrupt cell membrane. Consequently, the cell metabolisms may come to a halt and ultimately cell death to occur [33]. It seems ion channel or large pore formation (due to the lack of cholesterol in microbial cell membranes especially bacteria) within pathogen cell membrane is the key mechanisms of the most AMPs [34]. AMPs with the hydrophilic and hydrophobic tails manage their way through the phospholipids bilayer membranes of pathogens, leading to excess membrane permeation [35].

## CONCLUSION

Finally, it is clear from the present study that *Chrysomya megacephala* may be also useful in maggot therapy as *Lucilia sericata*. This also may reinforce the study of other maggots and their validity for larval therapy due to the presence of a lot of proteases in these maggots.

## ACKNOWLEDGEMENT

Deep thanks to Mycology Center at Al-AZHar University for their cooperation in Carrying this Work.

## REFERENCES

- Gupta A 2008: A review of the use of maggots in wound therapy. *Ann Plast Surg*, 60:224-227.
- Turkmen A, Graham K, Mc Grouther DA (2009): Therapeutic applications of the larvae for wound debridement. *Journal of Plastic, Reconstructive and Aesthetic Surgery*, 63:184-188.
- Chan DC, Fong DH, Leung JY, Patil NG and Leung GK. (2007). Maggot debridement therapy in chronic wound care. *Hong Kong Med J*, 13: 382-386.
- Zacur H and Kirsner RS. (2002). Debridement: rationale and therapeutic options. *Wounds* 14. Published online in September. ONLINE REF.
- Simmons SW (1935): A bactericidal principle in excretions of surgical maggots which destroys important etiological agents of pyogenic infections. *Bacteriol*, 30:253-267.
- Jaklič, D.; Lapanje, A.; Zupančič, K.; Smrke, D.; Gunde-Cimerman, N (2008). Selective antimicrobial activity of maggots against pathogenic bacteria. *J. Med. Microbiol.*, 57, 617–625.
- Daeschlein G, Mumcuoglu KY, Assadian O, Hoffmeister B, Kramer A, (2007): In vitro antibacterial activity of *Lucilia sericata* maggot secretions. *Skin Pharmacology and Physiology*, 20:112-115.
- Arora S, Lim CS, Baptista C (2010): Antibacterial activity of *Lucilia cuprinamaggot* extracts and its extraction techniques. *International Journal of Integrative Biology*, 9:43-48.
- Andersen AS, Sandvang D, Schnorr KM, Kruse T, Neve S, Joergensen B, Karlsmark T, Krogfelt KA (2010): A novel approach to the antimicrobial activity of maggot debridement therapy. *J Antimicrob Chemother* 65:1646-1654.
- Sukontason, K., Bunchoo, B. Khantawa, S.Piangjai, K. Sukontason, R. Methanitikorn and Y. Rongsriyam, (2000). Mechanical carrier of bacterial enteric pathogens by *Chrysomya megacephala* (Diptera: Calliphoridae) in Chiang Mai, Thailand. *Southeast Asian J Trop Med Public Health* 31(Suppl 1):157-161.
- Kumarasinghe, S.P.W., N.D. Karunaweera and R.L. Ihalamull (2000). A study of cutaneous myiasis in Sri Lanka. *Int J Dermatol* 39:689-694.
- Sukontason, K.L., P. Narongchai, K.Sukontason, R. Methanitikorn and S.Piangjai, (2005). Forensically important fly maggots in a floating corpse: the first case report in Thailand. *J Med Assoc Thai* 88:1458-1461.
- El-Ebiarie AS and Taha N (2012). Molecular characterization of serine proteases from both first and third larval instars of *Chrysomya megacephala*. *Life Science Journal*; 9(3):2086-2093.

14. Taha M. N. (2015). Activity of two serine proteases and anti-bacterial studies on both excretory/secretory products and whole body homogenates of third instar larvae of *Chrysomya megacephala*. International Journal of Advanced Research (2015), Volume 3, Issue 4, 451-459.
15. Gabre R. M., Fatma K. A., Hsin C (2005). Life table of *Chrysomya megacephala*. *Acta Oecologica* 27, 179-183.
16. Zumpt, F (1965). Myiasis in man and animals in the old-world. Butter Worths, London.
17. Bexfield A, Nigam Y, Thomas S, Ratcliffe NA (2004). Detection and partial characterisation of two antibacterial factors from the excretions/secretions of the medicinal maggot *Lucilia sericata* and their activity against methicillin-resistant *Staphylococcus aureus* (MRSA). *Microbes and Infection*, 6:1297-1304.
18. Merril CR (1990) "Gel-staining techniques" In Deutscher MP (Ed) Guide to protein purification. Methods in Enzymology volume 182. Academic press Inc.
19. Scott AC. (1989). Laboratory control of antimicrobial therapy. In: Collee JG *et al.* eds. *Practical Medical Microbiology*, 13th edn. Edinburgh: Churchill Livingstone: 161-181.
20. Bexfield, A.; Bond, A.E.; Roberts, E.C.; Dudley, E.; Nigam, Y.; Thomas, S.; Newton, R.P.; Ratcliffe, N.A. (2008). The antibacterial activity against MRSA strains and other bacteria of a <500 Da fraction of excretions/secretions of *Lucilia sericata* (Diptera: Calliphoridae). *Microbes Infect.* 10, 325-333.
21. Kerridge, A.; Lappin-Scott, H.; Stevens, J.R (2005). Antibacterial properties of larval secretions of the blowfly, *Lucilia sericata*. *Med. Vet. Entomol.*, 19, 333-337.
22. Jana Bohova, Juraj Majtan, Viktor Majtan, and Peter Takac (2014). Selective Antibiofilm Effects of *Lucilia sericata* Larvae Secretions/Excretions against Wound Pathogens. Hindawi Publishing Corporation Evidence-Based Complementary and Alternative Medicine Volume, Article ID 857360, 9 pages <http://dx.doi.org/10.1155/2014/857360>.
23. Lehane, M.J.; Aksoy, S.; Levashina, E. (2004). Immune responses and parasite transmission in blood-feeding insects. *Trends Parasitol.*, 20, 433-439.
24. Mak, P.; Zdybicka-Barabas, A.; Cytrynska, M. (2010). A different repertoire of *Galleriamellonella* antimicrobial peptides in larvae challenged with bacteria and fungi. *Dev. Comp. Immunol.*, 34, 1129-1136.
25. Hwang, J.-S.; Lee, J.; Kim, Y.-T.; Bang, H.-S.; Yun, E.-Y.; Kim, S.-R.; Suh, H.-J.; Kang, B.-R.; Nam, S.-H.; Joen, J.; Kim, I.; Lee, D.G. (2009). Isolation and characterization of a defensin-like peptide (Coprinsin) from the dung beetle, *Copris tripartitus*. *Int. J. Pept.*, doi:10.1155/2009/136284.
26. Song, K.-J.; Park, B.-R.; Kim, S.; Park, K. (2010). Molecular characterization of anionic defensin-like peptide in immune response of silkworm, *Bombyx mori* L. (Lepidoptera). *Genes. Genom.*, 32, 447-453.
27. Cunliffe, R.N.; Mahida, Y.R. (2004). Expression and regulation of antimicrobial peptides in the gastrointestinal tract. *J. Leukoc. Biol.*, 75, 49-58.
28. Rayaprolu, S.; Wang, Y.; Kanost, M.R.; Hartson, S.; Jiang, H. (2010). Functional analysis of four processing products from multiple precursors encoded by a leucocin-related gene from *Manduca sexta*. *Dev. Comp. Immunol.*, 34, 638-647.
29. Reddy, K.; Yedery, R.; Aranha, C. (2004). Antimicrobial peptides: premises and promises. *Int. J. Antimicrob. Ag.*, 24, 536-547.
30. Vizioli, J.; Salzet, M. (2002). Antimicrobial peptides from animals: focus on invertebrates. *Trends Pharmacol. Sci.*, 23, 494-496.
31. Mookherjee, N.; Hancock, R.E. (2007). Cationic host defence peptides: innate immune regulatory peptides as a novel approach for treating infections. *Cell. Mol. Life. Sci.*, 64, 922-933.
32. Papagianni, M. (2003). Ribosomally synthesized peptides with antimicrobial properties: biosynthesis, structure, function, and applications. *Biotechnol. Adv.*, 21, 465-499.
33. Liang, J.F.; Kim, S.C. (1999). Not only the nature of peptide but also the characteristics of cell membrane determine the antimicrobial mechanism of a peptide. *J. Pept. Res.*, 53, 518-522.
34. Melo, M.N.; Ferre, R.; Castanho, M.A. (2009). Antimicrobial peptides: linking partition, activity and high membrane-bound concentrations. *Nat. Rev. Microbiol.*, 7, 245-250.
35. Toke, O. (2005). Antimicrobial peptides: new candidates in the fight against bacterial infections. *Biopolymers*, 80, 717-735.

#### CITATION OF THIS ARTICLE

Nancy Taha Mohamed. Fractionation and Purification of Bioactive Peptides In Excretory/Secretory Products Of Third Instar Larvae Of *Chrysomya Megacephala* (Calliphoridae: Diptera). Bull. Env. Pharmacol. Life Sci., Vol 4 [10] September 2015: 69-74